

COWLES FOUNDATION FOR RESEARCH IN ECONOMICS

AT YALE UNIVERSITY

Box 2125, Yale Station
New Haven, Connecticut 06520

COWLES FOUNDATION DISCUSSION PAPER NO. 637

Note: Cowles Foundation Discussion Papers are preliminary materials circulated to stimulate discussion and critical comment. Requests for single copies of a Paper will be filled by the Cowles Foundation within the limits of the supply. References in publications to Discussion Papers (other than mere acknowledgment by a writer that he has access to such unpublished material) should be cleared with the author to protect the tentative character of these papers.

THE DISTRIBUTION OF MATRIX QUOTIENTS

by

Peter C. B. Phillips
Yale University

July 1982

THE DISTRIBUTION OF MATRIX QUOTIENTS

by

P. C. B. Phillips

Cowles Foundation
Yale University

0. ABSTRACT

Cramér's inversion formula for the distribution of a quotient is generalized to matrix variates and applied to give an alternative derivation of the matrix t -distribution.

KEYWORDS. Matrix variates, characteristic functions, matrix t -distribution

AMS CLASSIFICATION. Primary 62E15, Secondary 60E10

1. INTRODUCTION

Useful inversion formulae that apply for scalar ratios of random variates and proceed from the joint characteristic function of the component variates have been known for some time. In particular, if the scalar random variate $\eta \geq 0$ and has a finite mean Cramér [1] and Geary [3] give the following formula for the density of the ratio $\zeta = \xi/\eta$:

$$(1) \quad f(z) = \frac{1}{2\pi i} \int_{-\infty}^{\infty} \left[\frac{\partial \phi(s_1, s_2)}{\partial s_2} \right]_{s_2 = -zs_1} ds_1$$

where $\phi(s_1, s_2)$ is the joint characteristic function of (ξ, η) . Gurland [4] generalized (1) by considering the multidimensional case of a vector of ratios and relaxed the requirements that η necessarily be positive or have a finite mean (by using principal values in the integrals that define the inversions).

Closely related statistics that take the form of matrix quotients arise frequently in multivariate analysis. A common situation (leading, for example, to the matrix t-distribution) is the following. Let A be a positive definite $n \times n$ matrix variate partitioned as

$$(2) \quad A = \begin{bmatrix} & \ell & & k \\ A_{11} & & A_{21} & \\ & & & \\ A_{21} & & A_{22} & \\ & & & k \end{bmatrix} \begin{matrix} \ell \\ k \end{matrix}, \quad \ell+k = n$$

where A_{22} is a square submatrix of order k ; and suppose interest centers on the distribution of the quotient $X = A_{22}^{-1} A_{21}$. (For example, when A is central Wishart with degrees of freedom $T > k$, X is a regression

coefficient matrix for multivariate normal samples and is known to have a matrix t-distribution [2], [6], [7]). Corresponding to the matrix $F = (f_{ij})_{n \times n}$ we define the matrix ${}_{\eta}F = (\eta_{ij} f_{ij})$ where $\eta_{ij} = 1, 1/2$ for $i = j, i \neq j$ respectively. We denote the joint characteristic function of A by $\phi^*(F_{\eta}) = E\{\text{etr}(i_{\eta}FA)\}$. Partitioning F and ${}_{\eta}F$ conformably with A in (2), we define

$$(3) \quad \phi(F_{21}, {}_{\eta}F_{22}) = E\{\text{etr}(iF_{21}A'_{21} + i_{\eta}F_{22}A_{22})\} = E\{\text{etr}(2i_{\eta}F_{21}A'_{21} + i_{\eta}F_{22}A_{22})\} \\ = [\phi^*({}_{\eta}F)]_{\eta^{F_{11}}=0},$$

which is the joint characteristic function of the distinct elements of (A_{21}, A_{22}) . With this notation we develop an inversion formula for the joint density of the matrix quotient $X = A_{22}^{-1}A_{21}$ which generalizes (1) above.

THEOREM. *Suppose the joint density function $f(A_{21}, A_{22})$ of (A_{21}, A_{22}) exists everywhere and A_{22} is a positive definite matrix. Then, if $E(\det A_{22})^{\ell}$ exists, the density function of $X = A_{22}^{-1}A_{21}$ is given by*

$$(4) \quad f(X) = \left(\frac{1}{2\pi i}\right)^{k\ell} \int_{\mathbf{R}^{k\ell}} [D_{22}^{\ell} \phi(F_{21}, -(XF'_{21} + F_{21}X')/2)] dF_{21}$$

where D_{22} is the differential operator $\det(\partial/\partial {}_{\eta}F_{22})$.

2. PROOF OF THE THEOREM

By direct transformation of $(A_{21}, A_{22}) \rightarrow (X, A_{22})$ we deduce that

$$(5) \quad f(X) = \int_{A_{22} > 0} f(A_{22}X, A_{22}) (\det A_{22})^{\ell} dA_{22}.$$

We observe that the joint density

$$(6) \quad f^*(A_{21}, A_{22}) = [E(\det A_{22})^\ell]^{-1} (\det A_{22})^\ell f(A_{21}, A_{22})$$

defines a new distribution whose characteristic function

$E\{\text{etr}(iF_{21}'A_{21}' + i_{\eta}F_{22}'A_{22}')\}$ is given by

$$(7) \quad [E(\det A_{22})^\ell]^{-1} \int D_{22}^\ell \text{etr}(iF_{21}'A_{21}' + i_{\eta}F_{22}'A_{22}') f(A_{21}, A_{22}) dA_{21} dA_{22} \\ = [E(\det A_{22})^\ell]^{-1} D_{22}^\ell \phi(F_{21}, {}_{\eta}F_{22})$$

where the absolute convergence of the integral allows us to interchange the order of integration and differentiation.

Now consider the distribution of the matrix variate $W = A_{21} - A_{22}X$ given X where the joint distribution of (A_{21}, A_{22}) is defined by (6). The density of W is

$$(8) \quad f(W) = \int_{A_{22} > 0} f^*(W + A_{22}X, A_{22}) dA_{22} .$$

From (5) and (6) we see that $f(W)$ reduces to $[E(\det A_{22})^\ell]^{-1} f(X)$ when $W = 0$. We further note that the characteristic function of W is obtained by setting ${}_{\eta}F_{22} = -\frac{1}{2}(XF_{21}' + F_{21}X')$ in (7), that is

$$(9) \quad [E(\det A_{22})^\ell]^{-1} D_{22}^\ell \phi(F_{21}, {}_{\eta}F_{22}) \Big|_{{}_{\eta}F_{22} = -\frac{1}{2}(XF_{21}' + F_{21}X')} .$$

The required formula (4) for the density function $f(X)$ now follows from inversion of the characteristic function (9) and from taking its value at $W = 0$.

3. APPLICATION TO THE MATRIX t-DISTRIBUTION

Consider the canonical case of a central Wishart matrix A with degrees of freedom T and covariance matrix I_n . The joint characteristic function of A is

$$\phi^*(F_\eta) = [\det(I - 2iF_\eta)]^{-T/2}$$

and simple manipulations yield

$$(10) \quad \phi(F_{21}, {}_\eta F_{22}) = [\det(I - 2i({}_\eta F_{22} + F_{21}F'_{21}))]^{-T/2}.$$

Moreover,

$$(11) \quad D_{22}^\ell \phi(F_{21}, {}_\eta F_{22}) \propto [\det(I - 2i({}_\eta F_{22} + F_{21}F'_{21}))]^{-T/2-\ell}$$

(see, for example, [5] pp. 479-480). Substitution of (11) in (4) leads to the following expression for the density function of $X = A_{22}^{-1}A_{21}$:

$$(12) \quad f(X) \propto \int_{\mathbf{R}^{k\ell}} [\det\{I - i(XF'_{21} + F_{21}X') + F_{21}F'_{21}\}]^{-\frac{T}{2}-\ell} dF_{21} \\ = [\det(I+XX')]^{-\frac{T}{2}-\ell} \int_{\mathbf{R}^{k\ell}} [\det\{I - (I+XX')^{-\frac{1}{2}}(X+iF_{21})(X+iF_{21})'(I+XX')^{-\frac{1}{2}}\}]^{-\frac{T}{2}-\ell} dF_{21}.$$

We transform $F_{21} \rightarrow Z = (I+XX')^{-1/2}(F_{21} - iX)$ in the integral in (12).

This transformation has Jacobian $[\det(I+XX')]^{\ell/2}$ and we deduce that

$$(13) \quad f(X) \propto [\det(I+XX')]^{-(T+\ell)/2} \int_{\mathbf{R}^{k\ell}} [\det(I+ZZ')]^{-T/2-\ell} dZ.$$

The domain of integration in (13) can be taken to be $\mathbf{R}^{k\ell}$ as before since

the integrand is analytic in a strip of $C^{k\ell}$ that contains $R^{k\ell} - i(I+XX')^{-1/2}X$. It follows directly from (13) that the density function of $X = A_{22}^{-1}A_{21}$ is

$$f(X) = c[\det(I+XX')]^{-(T+\ell)/2}$$

where the constant in (14) takes the value

$$c = \pi^{-k\ell/2} \Gamma_{\ell} \left(\frac{1}{2}(T+\ell) \right) / \Gamma_{\ell} \left(\frac{1}{2}(T+\ell-k) \right)$$

as can be determined by elementary integration ([2] p. 512).

REFERENCES

- [1] Cramér, H. Mathematical Methods of Statistics, Princeton: Princeton University Press, 1946.
- [2] Dickey, J. M. "Matricvariate Generalizations of the Multivariate t Distribution and the Inverted Multivariate t Distribution," Annals of Mathematical Statistics (1967), 38, 511-518.
- [3] Geary, R. C. "Extension of a Theorem by Harold Cramér," Journal of the Royal Statistical Society (1944), 17, 56-57.
- [4] Gurland, J. "Inversion Formulae for the Distribution of Ratios," Annals of Mathematical Statistics (1948), 19, 228-237.
- [5] Herz, C. S., "Bessel Functions of Matrix Argument," Annals of Mathematics (1955), 61, 474-523.
- [6] Kabe, D. G. "On the Distribution of the Regression Coefficient Matrix of a Normal Distribution," Australian Journal of Statistics (1968), 10, 21-23.
- [7] Kshirsagar, A. M. "Some Extensions of the Multivariate t-distribution and the Multivariate Generalization of the Distribution of the Regression Coefficient," Proceedings of the Cambridge Philosophical Society (1960), 56, 80-85.